

Help Your Care Team Address Side Effects From **Pemazyre**



Know what to watch for while taking PEMAZYRE

PEMAZYRE is a prescription medicine that is used to treat adults with bile duct cancer (cholangiocarcinoma) that has spread or cannot be removed by surgery:

- who have already received a previous treatment, **and**
- whose tumor has a certain type of abnormal FGFR2 gene.

Your healthcare provider (HCP) will test your cancer for a certain type of abnormal FGFR2 gene and make sure that PEMAZYRE is right for you. It is not known if PEMAZYRE is safe and effective in children. PEMAZYRE is approved based on tumor response and duration of response. There are ongoing studies to show whether PEMAZYRE can improve survival or symptoms.

Your HCP may be able to offer ways to help manage side effects while you are taking PEMAZYRE.

Complete this checklist as needed between visits with your HCP and bring it with you if any of the boxes are checked.

PEMAZYRE may cause serious side effects, including:



EYE PROBLEMS

- Certain eye problems are common with PEMAZYRE but can also be serious
- Eye problems include: dry eye or inflamed eyes, inflamed cornea (front part of the eye), increased tears, and a disorder of the retina (an internal part of the eye)
- You will need to see an eye specialist for a complete eye exam before you begin treatment with PEMAZYRE, every 2 months for the first 6 months, and then every 3 months during treatment with PEMAZYRE
- You should use artificial tears or substitutes, hydrating or lubricating eye gels as needed, to help prevent or treat dry eyes

Tell your HCP right away if you develop any changes in your vision during treatment with PEMAZYRE including:

- blurred vision
- flashes of light
- black spots

! If you have these changes, report them to your HCP immediately. You may need to see an eye specialist right away.



HIGH PHOSPHATE LEVELS (HYPERPHOSPHATEMIA) AND BUILDUP OF MINERALS IN DIFFERENT TISSUES IN YOUR BODY

- Hyperphosphatemia is common with PEMAZYRE but can also be serious
- High levels of phosphate in your blood may lead to buildup of minerals such as calcium, in different tissues in your body
- Your healthcare provider will check your blood phosphate levels during treatment with PEMAZYRE
- Your healthcare provider may prescribe changes in your diet or phosphate lowering therapy, or change, interrupt or stop PEMAZYRE if needed

Tell your HCP right away if you develop:

- muscle cramps
- numbness or tingling around the mouth

! If you have these symptoms, report them to your HCP immediately. These can be symptoms of high phosphate levels.

The most common side effects of PEMAZYRE include:

- | | | |
|--|---|--|
| <input type="checkbox"/> hair loss | <input type="checkbox"/> constipation | <input type="checkbox"/> stomach-area (abdominal) pain |
| <input type="checkbox"/> diarrhea | <input type="checkbox"/> mouth sores | <input type="checkbox"/> low phosphate in blood* |
| <input type="checkbox"/> nails separate from the bed or poor formation of the nail | <input type="checkbox"/> dry eyes | <input type="checkbox"/> back pain |
| <input type="checkbox"/> feeling tired | <input type="checkbox"/> dry mouth | <input type="checkbox"/> dry skin |
| <input type="checkbox"/> change in sense of taste | <input type="checkbox"/> decrease in appetite | |
| <input type="checkbox"/> nausea | <input type="checkbox"/> vomiting | |
| | <input type="checkbox"/> joint pain | |

**Your HCP will measure this using lab tests.*

OTHER SIDE EFFECTS—If you experience any other side effects, please note them here and communicate them to your HCP.

Make sure to pay attention to signs and symptoms of potential side effects and communicate them to your HCP promptly. Your HCP may be able to offer ways to help manage side effects while you are taking PEMAZYRE.

These are not all the possible side effects of PEMAZYRE. Important Safety Information is shown on the back. Please visit PEMAZYRE.com to see the [Full Prescribing Information](#), including Patient Information, which includes a discussion of the risks associated with PEMAZYRE. For more information, ask your HCP or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

INDICATIONS AND USAGE

PEMAZYRE[®] is a prescription medicine that is used to treat adults with bile duct cancer (cholangiocarcinoma) that has spread or cannot be removed by surgery:

- who have already received a previous treatment, **and**
- whose tumor has a certain type of abnormal FGFR2 gene

Your healthcare provider will test your cancer for a certain type of abnormal FGFR2 gene and make sure that PEMAZYRE is right for you.

It is not known if PEMAZYRE is safe and effective in children.

PEMAZYRE is approved based on tumor response and duration of response. There are ongoing studies to show whether PEMAZYRE can improve survival or symptoms.

IMPORTANT SAFETY INFORMATION

Before you take PEMAZYRE, tell your healthcare provider about all of your medical conditions, including if you:

- have vision or eye problems
- have kidney problems
- have liver problems
- are pregnant or plan to become pregnant. PEMAZYRE can harm your unborn baby or cause loss of your pregnancy (miscarriage). You should not become pregnant during treatment with PEMAZYRE.

Females who can become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with PEMAZYRE.
- You should use an effective method of birth control during treatment and for 1 week after your last dose of PEMAZYRE. Talk to your healthcare provider about birth control methods that may be right for you.
- Tell your healthcare provider right away if you become pregnant or think that you may be pregnant.

Males with female partners who can become pregnant:

- You should use effective birth control when sexually active during treatment with PEMAZYRE and for 1 week after your last dose of PEMAZYRE.
- are breastfeeding or plan to breastfeed. It is not known if PEMAZYRE passes into your breast milk. Do not breastfeed during treatment and for 1 week after your last dose of PEMAZYRE.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of PEMAZYRE?

PEMAZYRE may cause serious side effects, including:

- **Eye problems.** Certain eye problems are common with PEMAZYRE but can also be serious. Eye problems include dry eye or inflamed eyes, inflamed cornea (front part of the eye), increased tears, and a disorder of the retina (an internal part of the eye). You will need to see an eye specialist for a complete eye exam before you begin treatment with PEMAZYRE, every 2 months for the first 6 months, and then every 3 months during treatment with PEMAZYRE.
 - You should use artificial tears or substitutes, hydrating or lubricating eye gels as needed, to help prevent or treat dry eyes.
 - **Tell your healthcare provider right away** if you develop any changes in your vision during treatment with PEMAZYRE, including: blurred vision, flashes of light, or see black spots. You may need to see an eye specialist right away.
- **High phosphate levels in your blood (hyperphosphatemia) and buildup of minerals in different tissues in your body.** Hyperphosphatemia is common with PEMAZYRE but can also be serious. High levels of phosphate in your blood may lead to buildup of minerals such as calcium, in different tissues in your body. Your healthcare provider will check your blood phosphate levels during treatment with PEMAZYRE.

- Your healthcare provider may prescribe changes in your diet or phosphate lowering therapy, or change, interrupt, or stop PEMAZYRE if needed.
- Tell your healthcare provider right away if you develop any muscle cramps, or numbness or tingling around your mouth.

The most common side effects of PEMAZYRE for cholangiocarcinoma include:

- | | |
|---|---------------------------------|
| • hair loss | • dry eyes |
| • diarrhea | • dry mouth |
| • nails separate from the bed or poor formation of the nail | • decrease in appetite |
| • feeling tired | • vomiting |
| • change in sense of taste | • joint pain |
| • nausea | • stomach-area (abdominal) pain |
| • constipation | • low phosphate in blood |
| • mouth sores | • back pain |
| | • dry skin |

These are not all the possible side effects of PEMAZYRE. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects.

You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Incyte Medical Information at 1-855-463-3463.

Keep PEMAZYRE and all medicines out of the reach of children.

General information about the safe and effective use of PEMAZYRE.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use PEMAZYRE for a condition for which it is not prescribed. Do not give PEMAZYRE to other people, even if they have the same symptoms you have. It may harm them. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information that is written for healthcare professionals.

Please see the [Full Prescribing Information](#) for PEMAZYRE.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

